

MULTITRANSITION NUCLEAR RELAXATION IN PRESENCE OF STATIC QUADROPOLAR INTERACTIONS

D. PETIT and J.-P. KORB

Laboratoire de Physique de la Matière Condensée, U.R.A. 1254 du Centre National de la Recherche Scientifique, Ecole Polytechnique, 91128 Palaiseau, France

1. Introduction

We have used the formalism of fictitious spin 1/2 to describe the nuclear relaxation of spins $I > 1/2$ in solids (1). We found a generalized macroscopic kinetic differential equation for any observable, expanded in a complete basis set of independent fictitious spin 1/2 operators, and whose time evolution is expressed in terms of a matrix of relaxation. The interest of such method was to allow a possible treatment of the relaxation whether for a single or multiple relaxation processes. Several examples were treated for dipolar and quadrupolar relaxations, including the case of a residual time averaged quadrupolar interaction (1).

However there exists some arguments against the choice of a basis of fictitious spin-1/2 operators. Several authors have proposed a description of the spin dynamics in term of an irreducible tensors basis (2, 3). Some have even claimed that the fictitious spin basis cannot reach the physical meaning of a studied phenomenon (3). We show on a particular case that one can obtain, without any calculation, a matrix of relaxation previously calculated (1) by using an energy level diagram which is a pictorial description of the fictitious spin operators.

It was also point out recently that the fictitious spin-1/2 basis presents some redundance (4). An alternative way of describing a spin 3/2 was proposed in term of a direct product of two Pauli matrices (Dirac basis (5)). We show that the fictitious spin-1/2 allows to build a basis without such redundance and the correspondance between the Dirac and fictitious spin-1/2 bases is given. We discuss finally some physical reasons to prefer either irreducible tensor or fictitious spins descriptions.

2. Relaxation in the case of multilevel systems

In Fig. 1 we have displayed three levels i, j, k of a multilevel system.




Fig. 1 Schematic description of the longitudinal relaxation of a three-level system, the different transitions are indicated as bold inclined lines, the deviations to global or partial equilibrium are also indicated.

The average of fictitious spin-1/2 operator $\langle I_z^{ij} \rangle$ describes the ij polarisation and the operators I_+^{ij} and I_-^{ij} describe the transitions $j \Rightarrow i$ and $i \Rightarrow j$, respectively. The dynamical variation of the polarisations are expressed by the following kinetic equations

$$\frac{d}{dt} \begin{pmatrix} \langle I_z^{ij} \rangle \\ \langle I_z^{jk} \rangle \end{pmatrix} = - \begin{pmatrix} \frac{1}{T_1^{ij,ij}} & \frac{1}{T_1^{ij,jk}} \\ 1 & 1 \\ \frac{1}{T_1^{jk,ij}} & \frac{1}{T_1^{jk,jk}} \end{pmatrix} \begin{pmatrix} \Delta \langle I_z^{ij} \rangle \\ \Delta \langle I_z^{jk} \rangle \end{pmatrix} \quad (1)$$

where the symbol Δ represents the deviation to equilibrium. Each $1/T_1^{ij,jk}$ rate, whose a general expression is given in ref(1), is a linear combination of spectral densities $J(\omega_{ij})$, ω_{ij} being the transition frequency. In Fig. (1) the inclined bold lines represent the equilibrium distributions of the global or partial ij

polarizations. The dashed vertical lines represent the initial distributions in case of saturation experiment. From Eq. (1), $\Delta\langle I_z^{ij} \rangle$ drives the dynamical variations of $\langle I_z^{ij} \rangle$ and $\langle I_z^{jk} \rangle$ through the relaxation rates $1/T_1^{ij,ij}$ and $1/T_1^{jk,ij}$, respectively. A transition has a positive contribution on such relaxation rates if it creates a variation of polarization which favors the return to equilibrium. For instance, the return to equilibrium $\Delta\langle I_z^{ij} \rangle$ let decrease the population of level j . Owing to the transition $j \Rightarrow k$, it thus results a negative evolution to equilibrium for the polarization $\langle I_z^{jk} \rangle$. In consequence the spectral density $J(\omega_{jk})$ will contribute negatively to $1/T_1^{jk,ij}$. Since the population j of $\langle I_z^{ij} \rangle$ is only involved here, this contribution occurs just once. Similarly the return to equilibrium $\Delta\langle I_z^{ij} \rangle$ increases the population i and the population k decreases due to the transition $k \Rightarrow i$. This drives effectively the polarization $\langle I_z^{jk} \rangle$ to equilibrium and yields a positive contribution of $J(\omega_{ik})$ to $1/T_1^{jk,ij}$. Let consider now the $1/T_1^{ij,ij}$ rate. The transition $i \Leftrightarrow j$ let decrease the population j and increases the population i during the return to equilibrium $\Delta\langle I_z^{ij} \rangle$. In consequence the spectral density $J(\omega_{ij})$ contributes positively twice to $1/T_1^{ij,ij}$. This gives finally :

$$\frac{1}{T_1^{jk,ij}} = -\alpha_{ij}J(\omega_{ij}) - \alpha_{jk}J(\omega_{jk}) + \alpha_{ik}J(\omega_{ik}),$$

$$\frac{1}{T_1^{ij,ij}} = +2\alpha_{ij}J(\omega_{ij}) + \alpha_{jk}J(\omega_{jk}) + \alpha_{ik}J(\omega_{ik}),$$

(2)

where the α_{ij} are some real positive coefficients immediately calculable in the interacting frame, using Eqs. (10 and 12) of Ref (1). For example in a quadrupolar relaxation process all the α_{ij} are equal.

3. Application to quadrupolar relaxation including a residual time averaged interaction

We have presented a comprehensive study of the relaxation of nuclear spins $I > 1/2$ in a previous paper (1) Here we apply the method described above to the quadrupolar relaxation of a nuclear spin $I=3/2$ in presence of a static or residual time averaged quadrupolar interaction H_Q .

The corresponding multilevel system is displayed on Fig. 2.




Fig.2 Schematic description of the longitudinal relaxation of a nuclear spin $I=3/2$ experiencing a high magnetic field and a residual quadrupolar interaction Ω_Q

The dynamical variation of the polarisations are expressed by the following equations :

$$\frac{d}{dt} \begin{pmatrix} \langle I_z^{14} \rangle \\ \langle I_z^{23} \rangle \\ \langle I_z^{12,34} \rangle \end{pmatrix} = - \begin{pmatrix} \frac{1}{T_1^{14,14}} & \frac{1}{T_1^{14,23}} & \frac{1}{T_1^{14,12-34}} \\ \frac{1}{T_1^{23,14}} & \frac{1}{T_1^{23,23}} & \frac{1}{T_1^{23,12-34}} \\ \frac{1}{T_1^{12-34,14}} & \frac{1}{T_1^{12-34,23}} & \frac{1}{T_1^{12-34,12-34}} \end{pmatrix} \begin{pmatrix} \Delta\langle I_z^{14} \rangle \\ \Delta\langle I_z^{23} \rangle \\ \Delta\langle I_z^{12,34} \rangle \end{pmatrix}$$

(3)

where the different relaxation rates are given in Table 1. All the coefficients of this Table can be found by a straightforward application of the diagrammatic method exposed in section 2.

It is worthwhile to make a comparison with the usual case of $\Omega_Q = 0$. Substituting $\Omega_Q = 0$ in Table 1 merges the columns (a - b) and (c - d). The longitudinal rates, noted in bold in Eq. (3), vanish. The resulting relaxation matrix is divided in two block matrices of dimension 2×2 and 1×1 , respectively. One can show that the former describes the dynamics of the Zeeman order and the latter deals with the quadrupolar order.

As already explained in ref. (1), the "bold" longitudinal coupling rates arise from two factors : (i) the presence of a residual time-averaged interaction (here $\Omega_Q \neq 0$); (ii) a drastic frequency dependence for the spectral density. These two factors are often induced by an anisotropic motion (low dimensionality,

restricted motion, ...). The immediate consequence of a motion of low symmetry is indeed to break the symmetry of the relaxation

$\frac{1}{T_{1Q}^{cd,ef}} = \frac{3}{2} C_Q^2 *$				
$\left[\begin{array}{l} a J_1(\omega_0 - \Omega_Q) + b J_1(\omega_0 + \Omega_Q) + \\ c J_2(2\omega_0 - \Omega_Q) + d J_2(2\omega_0 + \Omega_Q) \end{array} \right]$				
cd,ef	a	b	c	d
14,14	1	1	1	1
14,23	-1	-1	1	1
14,12-34	-1	1	-1	1
23,14=14,23				
23,23=14,14				
23,12-34	1	-1	-1	1
12-34,14	-2	2	-2	2
12-34,23	2	-2	-2	2
12-34,12-34	2	2	2	2

Table 1. Elements of the spin-lattice relaxation rates for the quadrupolar fluctuations in presence of a residual time-averaged quadrupolar splitting Ω_Q .

process itself. For instance in the case treated here, such a motion mixes together the dynamics of the Zeeman and the quadrupolar orders. So the eigendirections of the relaxation process in case of anisotropic motion are a mixture of the Zeeman and quadrupolar eigendirections for isotropic motions.(Eq. (27) of ref. (1).

4. Choice of the spin operators basis

We show here that the fictitious spin-1/2 basis does not present any redundancy and can be simply related to the other possible bases. This is detailed in Table 2 where we have given the 16 spin operators necessary to describe properly a spin 3/2 successively in the Dirac, fictitious spin 1/2 and irreducible tensors bases.

The presence of a redundancy in the fictitious spin 1/2 basis vanishes when choosing the spin operators A_p as follows :

$$A_p = \left\{ I_\alpha^{1j} \pm I_\alpha^{kl} \right\}, \alpha \in \{x, y, z\},$$

$$j \neq k \neq l \in \{2, 3, 4\}, p \in \{1, \dots, 15\} \text{ for } I = \frac{3}{2}$$

with $\text{Tr} \{ A_p A_q \} = \delta_{pq}$ (4)

The basis $\{A_p\}$ is completed by addition of the unity operator.

$\sigma_1 \otimes \sigma_1$ 1 T_0^3	$\sigma_1 \otimes \sigma_x \otimes x_2$ $2(I_x^{12} + I_x^{34})$ $-\frac{2\sqrt{3}}{5} T_1^3(a) + \frac{4}{3\sqrt{3}} T_1^3(s)$	$\sigma_1 \otimes \sigma_y \otimes y_2$ $2(I_y^{12} + I_y^{34})$ $\frac{2i\sqrt{3}}{5} T_1^3(s) + \frac{4i}{3\sqrt{3}} T_1^3(s)$	$\sigma_1 \otimes \sigma_z \otimes z_2$ $2(I_z^{12} + I_z^{34})$ $\frac{2}{5} T_0^3 + \frac{8}{3\sqrt{10}} T_0^3$
$\sigma_x \otimes \sigma_x \otimes x_1$ $2(I_x^{13} + I_x^{24})$ $\frac{2}{\sqrt{6}} T_2^3(s)$	$\sigma_x \otimes \sigma_x \otimes x_2$ $2(I_x^{14} + I_x^{23})$ $-\frac{2}{3} T_3^3(a) + \frac{2}{5} T_1^3(a)$ $+\frac{2}{\sqrt{15}} T_1^3(a)$	$\sigma_x \otimes \sigma_x \otimes y_2$ $2(I_y^{14} + I_y^{23})$ $\frac{2i}{3} T_3^3(s) + \frac{2i}{5} T_1^3(s)$ $+\frac{2i}{\sqrt{15}} T_1^3(s)$	$\sigma_x \otimes \sigma_x \otimes z_2$ $2(I_x^{13} - I_x^{24})$ $\frac{4}{3\sqrt{2}} T_2^3(s)$
$\sigma_y \otimes \sigma_x \otimes y_1$ $2(I_y^{13} + I_y^{24})$ $-\frac{2i}{\sqrt{6}} T_2^3(a)$	$\sigma_y \otimes \sigma_x \otimes y_2$ $2(I_y^{14} + I_y^{23})$ $\frac{2i}{3} T_3^3(s) + \frac{2i}{5} T_1^3(s)$ $-\frac{2i}{\sqrt{15}} T_1^3(s)$	$\sigma_y \otimes \sigma_y \otimes y_2$ $-2(I_x^{14} - I_x^{23})$ $\frac{2}{3} T_3^3(a) + \frac{2}{5} T_1^3(a)$ $+\frac{2}{\sqrt{15}} T_1^3(a)$	$\sigma_y \otimes \sigma_x \otimes z_2$ $2(I_y^{13} - I_y^{24})$ $-\frac{4i}{3\sqrt{2}} T_2^3(a)$
$\sigma_x \otimes \sigma_x \otimes z_1$ $2(I_x^{12} + I_x^{34})$ $\frac{4}{5} T_0^3 + \frac{4}{3\sqrt{10}} T_0^3$	$\sigma_x \otimes \sigma_x \otimes z_2$ $2(I_x^{12} - I_x^{34})$ $-\frac{2}{\sqrt{6}} T_2^3(a)$	$\sigma_x \otimes \sigma_x \otimes y_2$ $2(I_y^{12} - I_y^{34})$ $\frac{2i}{\sqrt{6}} T_2^3(s)$	$\sigma_x \otimes \sigma_x \otimes z_2$ $2(I_x^{12} - I_x^{34})$ $\frac{2}{\sqrt{6}} T_0^3$

Table 2 Correspondence between three bases describing a spin 3/2. The notations x, y, z corresponds to those of ref. (4). The definitions and properties of fictitious spin-1/2 can be found in ref. (6). The definitions of irreducible tensors $T(a)$ and $T(s)$ are given in ref. (3).

From Table 2, one notes that there is mainly two kinds of bases. The first kind is composed either of Dirac (product of cartesian spin operators) or fictitious spin-1/2 $\{A_p\}$ bases. The second kind involves the different possibilities to build an irreducible tensor basis (2, 3, 7).

The properties of the $\{A_p\}$ and product of cartesian spin operators bases are so close that in the description of a spin 3/2 with the $\{A_p\}$ basis we can use the same simple picture that the one previously proposed for the product of cartesian spin operators (8). This is precisely what we have displayed in Table 3. For exemple the term $z_1 x_2$ describes a x coherence of a spin "2" in antiphase with respect to spin "1". The corresponding A_p term describes the antiphase behavior of the one quantum coherence of the quadrupolar satellites 12 and 34.

Finally we discuss some physical reasons to prefer either irreducible or fictitious spins descriptions. It is difficult to define precisely the domain of application of one description or the other. However, it is often more adapted to use the irreducible tensor

description when the evolution operator itself can be seen as a simple rotation.

	in-phase one quantum x coherence quadrupolar satellite	in-phase one quantum y coherence quadrupolar satellite	polarization involving the Zeeman order
in-phase two quanta x coherence, two quanta quadrupolar satellite	"two-spin" coherence 3q. and central line of 1q. coherences	"two-spin" coherence 3q. and central line of 1q. coherences	antiphase two quanta x coherence, two quanta quadrupolar satellite
in-phase two quanta y coherence, two quanta quadrupolar satellite	"two-spin" coherence 3q. and central line of 1q. coherences	"two-spin" coherence 3q. and central line of 1q. coherences	antiphase two quanta y coherence, two quanta quadrupolar satellite
polarization involving the Zeeman order	antiphase one quantum x coherence quadrupolar satellite	antiphase one quantum y coherence quadrupolar satellite	longitudinal "two-spin" quadrupolar order

Table 3 Correspondence between the properties of the $\{A_p\}$ and product of cartesian spin operators bases. The place of each term is the same as in Table 2.

Effectively all the spin components are transformed in such a way that the global object "density matrix of spin 3/2" will not be distorted. For instance an experiment with "hard pulses" can be analyzed in term of rotations. On the other hand, when the evolution operator cannot be reduced to a simple rotation the global "density matrix of spin 3/2" is distorted. For exemple this occurs in experiments with selective pulses. Also the quadrupolar relaxation process in presence of static interactions treated above is another typical exemple. Effectively the mixing of the Zeeman and quadrupolar orders exhibits the break of the rotation symmetry. In this latter case the irreducible tensors would imply a linear combination of spherical tensors of different ranks. At least the fictitious spin-1/2 operators allows to follow directly the dynamics of the different level populations. This is always true because the picture coming from the $\{A_p\}$ basis is the more adapted to describe the populations and the transitions of a multilevel system.

References

- (1) D. Petit and J.-P. Korb, Phys. Rev. B 37, 5761 (1988).
- (2) B.C. Sanctuary, J. Chem. Phys. 64, 4352 (1976).

- (3) G.J. Bowden and W.D. Hutchinson, J. Magn. Res. 67, 403 (1986).
- (4) M. Goldman, Adv. Magn. Res., Symposium "High Resolution NMR in Solids", M.I.T. Jan. 1989.
- (5) G. Arfken, "The mathematical methods for physicists", Acad. Press, 3rd ed, 1985), Sect. 4.4, Table 4.1, P. 213.
- (6) S. Vega, J. Chem. Phys. 68, 5518 (1978).
- (7) D.M. Brink and G. R. Satchler, "Angular momentum", (Clarendon Press, Oxford, 1968).
- (8) (a) R. Ernst, G. Bodenhausen, and A. Wokaun, "Principles of nuclear magnetic resonance in one and two dimensions", (Oxford Science Publications, 1987), p. 173.
(b) M. Munowitz, "Coherence and NMR", (John Wiley & Sons, New York, 1988).